Author Index

Agabian, N., see Mottram, J.C., 115
Al-Yaman, F.M. and Knobloch, J.
Isolation and partial characterization of species-specific and cross-reactive antigens of *Echinococcus granulosus* cyst fluid, 101

Bakker, R.E., see Frenkel, M.J., 191
Beck, P., see Mason, P.J., 27
Ben-Ismail, R., see Smith, D.F., 213
Bender, K., see Hecker, H., 225
Betschart, B., see Hecker, H., 225
Bieber, D., see Zurita, M., 11
Black, S. and Vandeweerd, V.
Serum lipoproteins are required for multiplication of-*Trypanosoma brucei* brucei under axenic culture conditions, 65
Black, S.J., see Vandeweerd, V., 201
Boisvenue, R.J., see Shamansky, L.M., 73
Bordier, C., see Bouvier, J., 235
Bouvier, J., Bordier, C., Vogel, H., Reichelt, R. and Etges, R.
Characterization of the promastigote surface protease of *Leishmania* as a membrane-bound zinc endopeptidase, 235
Brown, K.N., see Snounou, G., 37
Brown, W.C., see Sugimoto, C., 159
Buening, G.M., see Hines, S.A., 1

Campbell, D.A.
A polyubiquitin gene from *Trypanosoma brucei (Short Communication)*, 147
Campbell, D.A., Suyama, D.A. and Simpson, L.
Genomic organisation of nuclear $tRNA^{Gly}$ and $tRNA^{Leu}$ genes in *Trypanosoma brucei*, 257
Conrad, P.A., see Sugimoto, C., 159
Cooper, D., see Kingston, I.B., 137
Cox, G.N., see Shamansky, L.M., 73

Da Cunha e Silva, N.L., Hassón-Voloch, A. and De Souza, W.
Isolation and characterization of a highly purified flagellar membrane fraction from trypanosomatids, 129
De Souza, W., see Da Cunha e Silva, N.L., 129
Dolan, T.T., see Sugimoto, C., 159
Donelson, J.E., see Engman, D.M., 285
Dore, E., see Pace, T., 109
Dragon, E.A., see Engman, D.M., 285

Edwards, M.R., Gilroy, F.V., Jimenez, B.M. and O'Sullivan, W.J.
Alanine is a major end product of metabolism by *Giardia lamblia:* a proton nuclear magnetic resonance study, 19
Engman, D.M., Sias, S.R., Gabe, J.D., Donelson, J.E. and Dragon, E.A.
Comparison of *HSP70* genes from two strains of *Trypanosoma cruzi (Short Communication)*, 285
Etges, R., see Bouvier, J., 235

Frenkel, M.J., Savin, K.W., Bakker, R.E. and Ward, C.W.
Characterization of cDNA clones coding from muscle tropomyosin of the nematode *Trichostrongylus colubriformis*, 191

Gabe, J.D., see Engman, D.M., 285
Gilroy, F.V., see Edwards, M.R., 19
Goddeeris, B.M., see Sugimoto, C., 159
Gramiccia, M., see Smith, D.F., 213

Hall, R., see Mason, P.J., 27
Haralambous, A., see Lightowlers, M.W., 171
Hassón-Voloch, A., see Da Cunha e Silva, N.L., 129
Hecker, H., Bender, K., Betschart, B. and Modespacher, U.-P.
Instability of the nuclear chromatin of procyclic *Trypanosoma brucei brucei*, 225
Hernández, F., see Rodríguez, M.A., 87
Hines, S.A., McElwain, T.F., Buening, G.M. and Palmer, G.H.
Molecular characterization of *Babesia bovis* merozoite surface proteins bearing epitopes immunodominant in protected cattle, 1
Hudson, L., see Takle, G.B., 57
Hundt, E., see Knapp, B., 47

Jarra, W., see Snounou, G., 37
Jimenez, B.M., see Edwards, M.R., 19

Karess, R.E., see Medina-Acosta, E., 263
Kemp, D.J., see Marshall, V.M., 281
Kingston, I.B., Wainwright, S.M. and Cooper, D., Comparison of collagen gene sequences in *Ascaris suum* and *Caenorhabditis elegans*, 137
Knapp, B., Hundt, E. and Küpper, H.A.
A new blood stage antigen of *Plasmodium falciparum* transported to the erythrocyte surface, 47
Knobloch, J., see Al-Yaman, F.M., 101
Küpper, H.A., see Knapp, B., 47

Le Page, R.W.F., see Tait, A., 247
Lew, A.M., see Marshall, V.M., 281
Lightowlers, M.W., Liu, D., Haralambous, A. and Rickard, M.D.
Subunit composition and specificity of the major cyst fluid antigens of *Echinococcus granulosus*, 171
Liu, D., see Lightowlers, M.W., 171
Lockyer, M.J., Marsh, K. and Newbold, C.I.
Wild isolates of *Plasmodium falciparum* show extensive polymorphism in T-cell epitopes of the circumsporozoite protein, 275

Mansour, T.E., see Zurita, M., 11
Marsh, K., see Lockyer, M.J., 275
Marshall, V.M., Peterson, M.G., Lew, A.M. and Kemp, D.J.
Structure of the apical membrane antigen I (AMA-1) of

Plasmodium chabaudi (Short Communication), 281
Mason, P.J., Shiels, B.R., Tait, A., Beck, P. and Hall, R.
Sequence and expression of a gene from *Theileria annulata* coding for a 70-kilodalton heat-shock protein
McElwain, T.F., see Hines, S.A., 1
Medina-Acosta, E., Karess, R.E., Schwartz, H. and Russell, D.G.
The promastigote surface protease (gp63) of *Leishmania* is expressed but differentially processed and localized in the amastigote stage, 263
Mertens, E., Van Schaftingen, E. and Müller, M.
Presence of a fructose-2,6-bisphosphate-insensitive pyrophosphate: fructose-6-phosphate phosphotransferase in the anaerobic protozoa *Tritrichomonas foetus*, *Trichomonas vaginalis* and *Isotricha prostoma*, 183
Modespacher, U.-P., see Hecker, H., 225
Mottram, J.C., Murphy, W.J. and Agabian, N., A transcriptional analysis of the *Trypanosoma brucei hsp83* gene cluster, 115
Mons, B., see Pace, T., 109
Murphy, W.J., see Mottram, J.C., 115
Mutharia, L.M., see Sugimoto, C., 159
Müller, M., see Mertens, E., 183

Newbold, C.I., see Lockyer, M.J., 275
Nicholls, S.C., see Takle, G.B., 57

O'Sullivan, W.J., see Edwards, M.R., 19
Orozco, E., see Rodriguez, M.A., 87

Pace, T., Ponzi, M., Dore, E., Scotti, R. and Mons, B.
Presence of contaminating mitochondrial DNA from host reticulocytes in experimental infections of *Plasmodium berghei*, 109
Palmer, G.H., see Hines, S.A., 1
Pearson, T.W., see Sugimoto, C., 159
Peterson, M.G., see Marshall, V.M., 281
Ponzi, M., see Pace, T., 109
Pratt, D., see Shamansky, L.M., 73

Ray, D.S., see Sheline, C., 151
Ready, P.D., see Smith, D.F., 213
Reichelt, R., see Bouvier, J., 235
Rickard, M.D., see Lightowlers, M.W., 171
Rodríguez, M.A., Hernández, F., Santos, L., Valdez, A. and Orozco, E.
Entamoeba histolytica: molecules involved in the target cell-parasite relationship, 87
Russell, D.G., see Medina-Acosta, E., 263

Santos, L., see Rodríguez, M.A., 87
Savin, K.W., see Frenkel, M.J., 191
Schwartz, H., see Medina-Acosta, E., 263
Scotti, R., see Pace, T., 109
Searle, S., see Smith, D.F., 213
Shamansky, L.M., Pratt, D., Boisvenue, R.J. and Cox, G.N.
Cuticle collagen genes of *Haemonchus contortus* and *Caenorhabditis elegans* are highly conserved, 73
Sheline, C. and Ray, D.S.
Specific discontinuities in *Leishmania tarentolae* minicircles map within universally conserved sequence blocks, 151
Shiels, B.R., see Mason, P.J., 27
Sias, S.R., see Engman, D.M., 285
Simpson, L., see Campbell, D.A., 257
Smith, D.F., Searle, S., Ready, P.D., Gramiccia, M. and Ben-Ismail, R.
A kinetoplast DNA probe diagnostic for *Leishmania major*: sequence homologies between regions of *Leishmania* minicircles, 213
Snary, D., see Takle, G.B., 57
Snounou, G., Jarra, W., Viriyakosol, S., Wood, J.C. and Brown, K.N.
Use of a DNA probe to analyse the dynamics of infection with rodent malaria prasites confirms that parasite clearance during crisis is predominantly strain- and species-specific, 37
Sugimoto, C., Mutharia, L.M., Conrad, P.A., Dolan, T.T., Brown, W.C., Goddeeris, B.M. and Pearson, T.W.
Protein changes in bovine lymphoblastoid cells induced by infection with the intracellular parasite *Theileria parva*, 159
Suyama, D.A., see Campbell, D.A., 257

Tait, A., Turner, C.M.R., Le Page, R.W.F. and Wells, J.M.
Genetic evidence that metacyclic forms of *Trypanosoma brucei* are diploid, 247
Tait, A., see Mason, P.J., 27
Takle, G.B., Young, A., Snary, D., Hudson, L. and Nicholls, S.C.
Cloning and expression of a trypomastigote-specific 85-kilodalton surface antigen gene from *Trypanosoma cruzi*, 57
Turner, C.M.R., see Tait, A., 247

Valdez, A., see Rodríguez, M.A., 87
Van Schaftingen, E., see Mertens, E., 183
Vandeweerd, V. and Black, S.J.
Serum lipoprotein and *Trypanosoma brucei brucei* interactions in vitro, 201
Vandeweerd, V., see Black, S., 65
Viriyakosol, S., see Snounou, G., 37
Vogel, H., see Bouvier, J., 235

Wainwright, S.M., see Kingston, I.B., 137
Ward, C.W., see Frenkel, M.J., 191
Wells, J.M., see Tait, A., 247
Wood, J.C., see Snounou, G., 37
Wong, S., see Campbell, D.A., 147

Young, A., see Takle, G.B., 57

Zurita, M., Bieber, D. and Mansour, T.E.
Identification, expression and in situ hybridization of an eggshell protein gene from *Fasciola hepatica*, 11

Subject Index

Adhesin, 87
Alanine, 19
Amastigote, 263
Amino acid metabolism, 19
Antigen 5, 171
Antigen B, 171
Apical membrane antigen, 281
Ascaris suum, 137

Babesia bovis, 1

Caenorhabditis elegans, 73, 137
Chromatin, 225
Circumsporozoite protein, 275
Collagen, 73
Collagen gene, 137
Conserved sequence block, 151
Crisis, 37
Cross-reactivity, 101
Cultivation, 73

Diagnostic probe, 213
DNA, 37
DNA sequence homology, 213

Echinococcus granulosus, 101, 171
Eggshell protein, 11
Eggshell protein expression, 11
Endopeptidase, 235
Energy metabolsim, 19
Entamoeba histolytica, 87, 183
Ethanol, 19

Fasciola hepatica, 11
Female genitalia of *F. hepatica*, 11
Flagellar membrane, 129
Flagellar pocket, 263
Flagellum, 129
Fructose-2,6-bisphosphate, 183

Gene cloning, 57
Gene structure, 73
Giardia lamblia, 19
Glycoprotein, 263

^{1}H-magnetic resonance spectroscopy, 19
Haemonchus contortus, 73
Heat-shock element, 115
Heat-shock induction, 27
Heat-shock protein, 27
Heat-shock protein gene (Hsp83), 115
High-density lipoprotein, 65
Histone, 225
HSP70 gene, 285
Hybridisation, 37
Hydatid cyst fluid, 101

Immunodiagnosis, 101
Immunoprecipitation, 171
Intergenic sequence, 115
Isotricha prostoma, 183

Leishmania, 235, 263
Leishmania major, 213
Leishmania tarentolae, 151
Lipid anchor, 1
Lipid uptake, 201
Lipoprotein, 201
Lipoprotein-lipid, 65
Low-density lipoprotein, 65

Malaria, 37
Meiosis, 247
Membrane protein, 235
Merozoite, 1
Metacyclic, 247
Metalloprotease, 235
Minicircle kDNA, 213
Mitochondrial DNA, 109
Multiplication in axenic cultures, 65
Muscle, 191

Nucleosome filament, 225

Phenylhydrazine treatment, 109
Phosphatidylinositol, 1
Phosphofructokinase, 183
Phosphorylcholine, 171
Plasmodium berghei, 37, 109
Plasmodium chabaudi, 37, 281
Plasmodium falciparum, 47, 275
Ploidy, 247
Polyubiquitin gene, 147
Protease, 235
Protein structure, 47
Psoralen cross-linking, 225
Purification, 101
Pyrophosphate, inorganic, 183

Reticulocyte maturation, 109
RNA polymerase III promoter, 257
Rodent, 37

Schizont, 159
Schizont protein, 47
Sequence homology, 73
Serological diagnosis, 171
Specific discontinuity, 151
Specificity, 101, 171
Surface antigen, 1
Surface antigen, 85 kDa, 57
Surface location, 47
Surface glycoprotein, 159
Surface protein, 87, 159

Synchronised infections, 109

T cell epitope, 275
T lymphoblast, 159
Theileria annulata, 27
Theileria parva, 159
Transfer RNA gene, 257
Trichomonas vaginalis, 183
Trichostrongylus colubriformis, 191
Triton X-114, 1
Tritrichomonas foetus, 183
Tropomyosin, 191
Trypanosoma brucei, 115, 147, 247, 257
Trypanosoma brucei brucei, 65, 201
Trypanosoma cruzi, 47, 285
Trypanosomatids, 129
Trypanosome, 225
Trypomastigote, 57
Two-dimensional gel electrophoresis, 159

Ultrastructure, 129

Zinc, 235